

**TRANSVERSE DOUBLE-SPIN ASYMMETRIES
FOR SMALL Q_T DRELL-YAN PAIR PRODUCTION
IN PP AND $P\bar{P}$ COLLISIONS**

H. KAWAMURA

Radiation Laboratory, RIKEN, Wako 351-0198, Japan

E-mail: hiroyuki@rarfexp.riken.jp

J. KODAIRA

Theory Division, KEK, Tsukuba 305-0801, Japan

E-mail: jiro.kodaira@kek.jp

K. TANAKA

Department of Physics, Juntendo University, Inba, Chiba 270-1695, Japan

E-mail: tanakak@sakura.juntendo.ac.jp

We discuss transverse double-spin asymmetries for dimuon production at small transverse-momentum Q_T in pp and $p\bar{p}$ collisions. All order resummation of large logarithms relevant in small Q_T region is performed at next-to-leading logarithmic (NLL) accuracy, and asymmetries at RHIC, J-PARC and GSI are calculated.

The transversity $\delta q(x)$, the distribution of transversely polarized quarks inside transversely polarized nucleon, is the last unknown distribution of nucleon at the leading twist.¹ It is not measurable in inclusive DIS due to its chiral-odd structure, and a number of experiments are underway to measure it through semi-inclusive processes. Transversely polarized Drell-Yan process (tDY) is another way to measure it, which could in principle provide us with clean information on the transversity. However, the actual feasibility of extracting $\delta q(x)$ from tDY data depends on the magnitude of transverse double-spin asymmetry: $\frac{\Delta_T d\sigma}{d\sigma} \equiv \frac{d\sigma^{\uparrow\uparrow} - d\sigma^{\uparrow\downarrow}}{d\sigma^{\uparrow\uparrow} + d\sigma^{\uparrow\downarrow}}$. At RHIC, the asymmetry is possibly small since (i) pp collision probes products of valence-quark and sea-antiquark distributions where the latter are supposed to be small, and (ii) the small- x rise of unpolarized sea-quark distributions enhance the denominator of the asymmetry at high energy.² On the other hand, possibilities of transversely polarized pp ($p\bar{p}$) experiments at J-PARC (GSI) with moderate energies are recently discussed, where larger asym-

metries in tDY are expected.³ In this paper, we explore the asymmetries in Q_T spectrum of Drell-Yan pair, especially in small Q_T region where the bulk of the lepton pair is produced.

In the small Q_T region, fixed-order perturbative calculation does not work well since there appear large logarithmic corrections at each order of perturbation theory as $\alpha_s^n \ln^{2n-1}(Q^2/Q_T^2)/Q_T^2$, $\alpha_s^n \ln^{2n-2}(Q^2/Q_T^2)/Q_T^2$, and so on, corresponding to LL, NLL, and higher contributions, respectively. These “recoil logs” have to be resummed to all orders in α_s to obtain reliable perturbative predictions. The resummation is carried out in the impact parameter b space, conjugate to the Q_T space, to take transverse-momentum conservation into account, and the resummed cross section is expressed as the Fourier transform back to the Q_T space.⁴ At the NLL, the resummed cross section of tDY, differential in invariant mass Q , transverse momentum Q_T and rapidity y of the lepton pair, and in azimuthal angle ϕ of one of the outgoing leptons is given by⁵ ($\sqrt{S}$ the CM energy of hadrons)

$$\frac{\Delta_T d\sigma^{\text{NLL}}}{dQ^2 dQ_T^2 dy d\phi} = \cos(2\phi) \frac{\alpha^2}{3 N_c S Q^2} \sum_i e_i^2 \int_0^\infty db \frac{b}{2} J_0(bQ_T) e^{S(b,Q)} \quad (1)$$

$$\times \left[(C_{qq} \otimes \delta q_i) \left(x_1^0, \frac{b_0^2}{b^2} \right) (C_{\bar{q}\bar{q}} \otimes \delta \bar{q}_i) \left(x_2^0, \frac{b_0^2}{b^2} \right) + (x_1^0 \leftrightarrow x_2^0) \right].$$

Here $J_0(bQ_T)$ is a Bessel function, $b_0 = 2e^{-\gamma_E}$ with γ_E the Euler constant, and the large logarithmic corrections are resummed into the Sudakov factor $e^{S(b,Q)}$ with $S(b,Q) = -\int_{b_0^2/b^2}^{Q^2} (d\kappa^2/\kappa^2) \{ \ln \frac{Q^2}{\kappa^2} A_q(\alpha_s(\kappa)) + B_q(\alpha_s(\kappa)) \}$. A_q , B_q and the coefficient functions $C_{qq}, C_{\bar{q}\bar{q}}$ are perturbatively calculable, and are found in Ref.⁵ up to the accuracy necessary for the NLL. $x_{1,2}^0 = (Q/\sqrt{S})e^{\pm y}$, and $\delta q_i(x, \mu^2)$ is the transversity of i -th flavour quark at the $\overline{\text{MS}}$ scale μ . The singularity in b -integration,⁴ due to the Landau pole in $\alpha_s(\kappa)$, is taken care of by “contour deformation method” introduced in the joint resummation;⁷ correspondingly, ^{4,6,7} we take non-perturbative effects into account by the replacement $e^{S(b,Q)} \rightarrow e^{S(b,Q) - g_{NP} b^2}$ in (1), with a non-perturbative parameter g_{NP} . We combine the resulting NLL cross section with the leading order (LO) cross section, which is of $\mathcal{O}(\alpha_s)$ and is obtained⁵ as QCD prediction at large Q_T ; the matching of the NLL formula with the corresponding component in the LO cross section is performed at intermediate Q_T following the formulation developed in Ref.⁶, to ensure no double counting for all Q_T , and we finally obtain the complete “NLL+LO” cross section $\Delta_T d\sigma/(dQ^2 dQ_T^2 dy d\phi)$, which has a uniform accuracy over the entire range of Q_T .⁵ Note that, also for the unpolarized DY, the corre-

sponding “NLL+LO” cross section $d\sigma/(dQ^2 dQ_T^2 dy d\phi)$ can be obtained in the same framework, utilizing the results in the literatures.^{10,4.}

We calculate the following transverse double-spin asymmetries:

$$A_{TT} = [\Delta_T d\sigma/dQ^2 dQ_T^2 dy d\phi] / [d\sigma/dQ^2 dQ_T^2 dy d\phi] . \quad (2)$$

As non-perturbative inputs, we use the same parton distributions as those used in Ref.²; in particular, for the numerator, we use a model of the transversity $\delta q(x, \mu^2)$ which saturates the Soffer bound as $\delta q(x, \mu_0^2) = [q(x, \mu_0^2) + \Delta q(x, \mu_0^2)]/2$ at low input scale $\mu_0 \sim 0.6$ GeV and is evolved to higher μ^2 with NLO DGLAP kernel.⁸ The non-perturbative parameter g_{NP} is taken to be the same value for both numerator and denominator of (2), and we use $g_{NP} \simeq 0.5$ GeV² as suggested by the result of Ref.⁹.




Figure 1. A_{TT} in pp collision. Left: $\sqrt{S} = 200$ GeV, $y = 2$, $\phi = 0$ and $Q = 3, 5, 8, 15$ GeV, corresponding to RHIC kinematics. Right: $\sqrt{S} = 10$ GeV, $y = \phi = 0$ and $Q = 2, 3, 4$ GeV, corresponding to J-PARC kinematics. $g_{NP} = 0.5$ GeV² in both cases.

A_{TT} in pp collision are shown as functions of Q_T in Fig. 1, where the left (right) panel is for RHIC (J-PARC) kinematics. $A_{TT} \gtrsim 10\%$ are obtained for J-PARC, where the parton distributions are probed at medium $x_{1,2}^0$ (see (1)). For the RHIC case, A_{TT} are less than 10%, and becomes smaller for smaller Q due to the small- x rise of unpolarized sea-distributions in the denominator of (2). The largest A_{TT} of 15-30% are obtained in $p\bar{p}$ collision at GSI kinematics as shown in Fig. 2, where A_{TT} are dominated by valence distributions at medium x . Integrating the numerator and denominator of (2) over Q_T^2 , our results for GSI kinematics reproduce the corresponding NLO asymmetries given by Barone et al.³.

In all cases of Figs. 1 and 2, A_{TT} have flat behavior in small Q_T region, although the numerator and denominator of (2) have strong Q_T dependence, respectively.⁵ We also note that A_{TT} are almost unaffected when




Figure 2. A_{TT} in $p\bar{p}$ collision at $\sqrt{S} = 14.5$ GeV, $y = \phi = 0$, and $Q = 2, 3, 4, 6$ GeV, corresponding to GSI kinematics, and with $g_{NP} = 0.5 \text{ GeV}^2$.

the parameter g_{NP} is varied in the range $g_{NP} = 0.3\text{--}0.8 \text{ GeV}^2$. These common features in Figs. 1 and 2 come from the dominance of soft gluon effects in small Q_T region, whose main part, i.e. the Sudakov factor of (1), is universal⁵ at the NLL level in both polarized and unpolarized channels.

Acknowledgments

We would like to thank Werner Vogelsang and Stefano Catani for valuable discussions. The work of J.K. and K.T. was supported by the Grant-in-Aid for Scientific Research Nos. C-16540255 and C-16540266.

References

1. J. P. Ralston and D. Soper, *Nucl. Phys.*, **B152**, 109 (1979).
2. O. Martin, et al., *Phys. Rev.* **D57**, 3084 (1998); *ibid.* **D60**, 117502 (1999).
3. H. Shimizu et al., *Phys. Rev.* **D71** (2005) 114007; H. Yokoya, in these proceedings; V. Barone et al., hep-ph/0512121; P. Lenisa in these proceedings.
4. J. C. Collins, D. Soper and G. Sterman, *Nucl. Phys.* **B250** 199 (1985).
5. H. Kawamura, J. Kodaira, H. Shimizu and K. Tanaka, *Prog. Theor. Phys.* **115**, 667 (2006).
6. G. Bozzi, S. Catani, D. de Florian and M. Grazzini, *Phys. Lett.* **B564** (2003) 65; *Nucl. Phys.* **B737** (2006) 73.
7. E. Laenen, G. Sterman and W. Vogelsang, *Phys. Rev.* **D63** (2001) 114018; S. Kulesza, G. Sterman and W. Vogelsang, *Phys. Rev.* **D66** (2002) 014001.
8. A. Hayashigaki, Y. Kanazawa and Y. Koike, *Phys. Rev.* **D56** 7350 (1997); S. Kumano and M. Miyama, *Phys. Rev.* **D56** 2504 (1997); W. Vogelsang, *Phys. Rev.* **D57** 1886 (1998).
9. A. Kulesza and W. J. Stirling, *JHEP* **0312** (2003) 056.
10. G. Altarelli et al., *Nucl. Phys.* **B246**, 12 (1984).